

Ground state energy of the modified Nambu–Goto string.

Leszek Hadasz[†]

Jagiellonian University, Institute of Physics
Reymonta 4, 30–059 Kraków, Poland

Abstract

We calculate, using zeta function regularization method, semiclassical energy of the Nambu–Goto string supplemented with the boundary, Gauss–Bonnet term in the action and discuss the tachionic ground state problem.

TP-JU 12/97
November 1997
hep-th/9711101

[†]E-mail: hadasz@thp1.if.uj.edu.pl

It seems to exist a common belief that some string representation of QCD would be very helpful in understanding such non-perturbative properties of quantum chromodynamics as the nature of the ground state or mechanism of confinement. The conjecture of existence of such (at least approximate) description is supported by a number of facts [1], to mention only the nature of the $1/N_c$ expansion [2], success of the dual models in description of Regge phenomenology, area confinement law found in the strong coupling lattice expansion [3] or the existence of flux-line solutions in confining gauge theories [4, 5] and the analytical results concerning two-dimensional QCD [6].

The natural zeroth order approximation in constructing the string description of QCD, suggested by the approaches mentioned above, is the Nambu-Goto string [7]. Unfortunately, when treated as a quantum system, it has many well known drawbacks [8, 9], which include the non-physical dimension of the space-time ($D=26$) or tachion and unwanted massless states in the spectrum.

In the free Nambu-Goto string model the squared mass of the ground state is (up to some positive factor with the corresponding dimension [8]) just the Casimir energy [11]. Therefore, revealing the sign of this energy we can demonstrate the absence or persistence of a tachion in the spectrum, indicating in this way the possibility of constructing a complete, consistent quantum theory for the string model under consideration or invalidating such a construction.

A model which has been extensively investigated along this line is the Nambu-Goto string with the point-like masses attached at the ends [12]. The results – although not decisive for the string propagating in the physical, four dimensional space-time – are encouraging.

In this letter we want to consider another modification: Nambu-Goto string with the boundary, Gauss-Bonnet term added (for details see [13, 14]), defined by the action:

$$S = \int d^2\xi \sqrt{-g} \left(-\gamma - \frac{\alpha}{2} R \right), \quad (1)$$

where

$$g_{ab} = \partial_a X_\mu \partial_b X^\mu, \quad a, b = \tau, \sigma,$$

$$g = \det(g_{ab}),$$

X_μ gives immersion of the two-dimensional string world-sheet parameterized by (τ, σ) into the four-dimensional Minkowski spacetime, γ and α are constants and R is the inner curvature of the string world-sheet.

The addition of the Gauss-Bonnet term is a rather natural construction in the context of the effective QCD string. The QCD string action should contain – apart from the X^μ fields – also infinitely many fields describing for instance the transverse shape of the chromoelectric flux joining the color sources. In constructing the effective string action one integrates over such a fields and this procedure inevitably leads to emergence of the intrinsic curvature term in the action functional. Of course, it is then only the first one out of the infinitely many terms with the growing number of derivatives.

In the four-dimensional space time (which we assume in this paper) all the classical solutions of the model defined by the action functional (1) can be obtained (for details see [8, 13]) by solving the complex Liouville equation [15]:

$$\ddot{\Phi} - \Phi'' = 2q^2 e^{\Phi}, \quad (2)$$

supplemented with the boundary conditions:

$$e^{\Phi} = -\frac{1}{q} \sqrt{\frac{\alpha}{\gamma}}, \quad \Im \Phi' = 0 \quad \text{for} \quad \sigma = \pm \frac{\pi}{2}. \quad (3)$$

The distinguished class of solutions of the Eqs. (2,3) consists of static (i.e. τ -independent) Liouville fields. This solutions are of the form:

$$\Phi_0 = \log \left(\frac{1}{q^2} \frac{\lambda^2}{\cos^2 \sigma} \right) + i\pi, \quad (4)$$

where the parameters of the model α, γ and the parameters of the solution λ, q are connected by the condition

$$\frac{\cos^2 \frac{\pi\lambda}{2}}{\lambda^2} = \frac{1}{q} \sqrt{\frac{\alpha}{\gamma}}, \quad (5)$$

following from Eq. (3). The Liouville field (4) corresponds to the string which rotates rigidly in a plane:

$$(X^\mu) = \frac{q}{\lambda^2} (\lambda\tau, \cos \lambda\tau \sin \lambda\sigma, \sin \lambda\tau \sin \lambda\sigma, 0). \quad (6)$$

In the paper [14] small oscillations around the solution (4) have been investigated. If we write the full Liouville field Φ in the form:

$$\Phi = \Phi_0 + \Phi_1, \quad (7)$$

and assume that Φ_1 is small, then we have:

$$\ddot{\Phi}_1 - \Phi_1'' = -V_\lambda(\sigma) \Phi_1, \quad (8)$$

together with the boundary conditions:

$$\Phi_1 = 0; \quad \Im \Phi_1' = 0 \quad \text{for} \quad \sigma = \pm \frac{\pi}{2}. \quad (9)$$

Here we have defined:

$$V_\lambda(\sigma) = \frac{2\lambda^2}{\cos^2 \lambda\sigma}. \quad (10)$$

The general solution of the Eqs. (8,9) is of the form:

$$\Phi_1(\tau, \sigma) = \sum_{n=1}^{\infty} D_n \cos(\omega_n \tau + \varphi_n) \left\{ \tan \lambda\sigma \cos \left(\omega_n \sigma + \frac{\pi n}{2} \right) - \frac{\omega_n}{\lambda} \sin \left(\omega_n \sigma + \frac{\pi n}{2} \right) \right\}, \quad (11)$$

where D_n and φ_n are arbitrary constants and the eigenfrequencies ω_n are roots of the equation:

$$\omega_n \tan \left[\frac{\pi(\omega_n + n)}{2} \right] = \lambda \tan \left(\frac{\pi\lambda}{2} \right), \quad (12)$$

for $n \geq 1$. The contribution to the string energy due to the fluctuation $\Phi_1(\tau, \sigma)$ is positive [14]. This shows the classical stability of the rigidly rotating string against small perturbations.

The constraints following from the reparametrization invariance of the action functional (1) reduce the number of the functional d.o.f. of the X^μ field by two. The boundary conditions, following from the higher derivative terms in the action, impose additional restriction which, in the case of considered small oscillations, (9), forces the Liouville field (11) to be real. This means that the string's functional d.o.f. are finally reduced to one. We can for instance describe the fluctuations of the string through the distance of the point with the coordinate σ from the center of rotation $\sigma = 0$, evaluated at some fixed laboratory time X^0 ,

$$l(\tau, \sigma) = \frac{q}{\lambda^2} \sin \lambda \sigma + \frac{2q}{\lambda^2} \sum_{n=1}^{\infty} D_n \left\{ \frac{\omega_n^+}{\omega_n^-} \cos \left(\omega_n^- \sigma + \frac{\pi n}{2} \right) + \frac{\omega_n^-}{\omega_n^+} \cos \left(\omega_n^+ \sigma + \frac{\pi n}{2} \right) \right\} \sin \left(\frac{\lambda \omega_n}{q} X^0 + \varphi_n \right) \quad (13)$$

where $\omega_n^\pm = \omega_n \pm \lambda$.

Formula (13) shows that oscillations of the weakly perturbed, rotating string can be expressed as a sum over non-interacting modes with the frequencies

$$\nu_n = \frac{\lambda \omega_n}{q}. \quad (14)$$

This suggests [16] that in the semiclassical (i.e. quadratic in the amplitudes of the oscillations) approximation the quantum string Hamiltonian can be written as

$$H = E_0 + \frac{1}{2} \sum_{n=1}^{\infty} \nu_n (b_n^\dagger b_n + b_n b_n^\dagger) = E_0 + \sum_{n=1}^{\infty} \nu_n b_n^\dagger b_n + \frac{1}{2} \sum_{n=1}^{\infty} \nu_n, \quad (15)$$

where

$$E_0 = \frac{\gamma q \pi}{2\lambda} \left(1 + \frac{\sin \pi \lambda}{\pi \lambda} \right) \quad (16)$$

is the classical string mass and the operator $b_n^\dagger$ (b_n) creates (annihilates) the n -th mode of the string oscillations.

The Hilbert space of the quantized string is now constructed as a Fock space generated by the creation operators $b_n^\dagger$ in the standard way [16]. The only term that needs some care is the last one in the formula (15).

To sum over mode frequencies which appear in the equation (15) is divergent and in order to give it a physical meaning we have to adopt some regularization scheme. We choose to work with the ζ function regularization [17] and define

$$E_C \stackrel{\text{def}}{=} \frac{1}{2} \lim_{s \rightarrow -1} \mu^{s+1} \zeta(s) \quad (17)$$

where, for $\Re s > 1$,

$$\zeta(s) = \sum_{n=1}^{\infty} \nu_n^{-s} \quad (18)$$

and the parameter μ with dimension of mass was introduced to ensure that the r.h.s. of the expression (17) has the dimension of energy for arbitrary complex s . The physically interesting value $s = -1$ is obtained from (18) through the analytic continuation.

In order to evaluate (17) let us first rewrite the equation (12) for ω_n in the equivalent form,

$$D(\lambda, \omega) \stackrel{\text{def}}{=} \sin(\pi\omega) \left[\eta^2 - \frac{1}{\omega^2} - \frac{2\eta}{\omega} \cot(\pi\omega) \right] = 0, \quad (19)$$

where $\eta^{-1} = \lambda \tan \frac{\pi\lambda}{2}$. Using the standard methods of contour integration in the complex plane we have

$$\sum_{n=1}^N \omega_n^{-s} = \frac{1}{2\pi i} \int_{C_1} dz z^{-s} \frac{d}{dz} \log D(\lambda, z), \quad (20)$$

where the integration contour C_1 (Fig. 1) surrounds N first zeroes of the function D .




Fig 1. The integration contours in the complex plane.

Let us now deform the integration contour C_1 into C_2 which consists of a semi circle of radius R , two line intervals on imaginary axis $[i\epsilon, iR] \cup [-i\epsilon, -iR]$ and a semi circle of radius ϵ . In the limit $R \rightarrow \infty$ the integral over the outer semi circle vanishes and we are left with the expression

$$\sum_{n=1}^{\infty} \omega_n^{-s} = \frac{1}{\pi} \sin\left(\frac{\pi s}{2}\right) \left\{ \frac{\epsilon^{-s}}{s} + \int_{\epsilon}^{\infty} dx x^{-s} [\pi \coth \pi x - \rho(\eta, x)] \right\}, \quad (21)$$

where

$$\rho(\eta, x) = \frac{2}{x} \frac{1 + \eta x \coth(\pi x) + \eta \pi x^2 \operatorname{cosech}^2(\pi x)}{1 + 2\eta x \coth(\pi x) + \eta^2 x^2}.$$

In order to be able to analytically continue r.h.s. of the equation (21) to the vicinity of $s = -1$ we subtract from the integrand its (uniform in η) expansion for $x \rightarrow \infty$,

$$\int_{\epsilon}^{\infty} dx x^{-s} [\pi \coth \pi x - \rho(\eta, x)] = \int_{\epsilon}^{\infty} dx x^{-s} \left[\pi \coth \pi x - \rho(\eta, x) - \pi + \frac{2}{x + \eta x^2} \right] +$$

(22)

$$+ \int_{\varepsilon}^{\infty} dx x^{-s} \left[\pi - \frac{2}{x + \eta x^2} \right].$$

Defining

$$F(\eta, s) = \int_{\varepsilon}^{\infty} dx \frac{x^{-s}}{x + \eta x^2}$$

we have (for $\Re s > 1$)

$$F(\eta, s) = \int_{\varepsilon}^{\infty} dx x^{-(s+1)} \left[1 - \eta \frac{x}{1 + \eta x} \right] = \frac{\varepsilon^{-s}}{s} - \eta F(\eta, s-1),$$

or equivalently

$$F(\eta, s) = \frac{1}{\eta} \left[\frac{\varepsilon^{-(s+1)}}{s+1} - F(\eta, s+1) \right].$$

This relations allows us to analytically continue the function $F(\eta, s)$ to the vicinity of $s = -1$ with the result

$$F(\eta, s) \stackrel{s \rightarrow -1}{=} \frac{1}{\eta} \frac{1}{s+1} + \mathcal{O}(s+1).$$

Analytic continuation of the integral

$$\int_{\varepsilon}^{\infty} dx \pi x^{-s}$$

gives zero and we finally arrive at the expression

$$\sum_{n=1}^{\infty} \omega_n^{-s} \stackrel{s \rightarrow -1}{=} \frac{2}{\pi \eta} \frac{1}{s+1} - \frac{1}{12} + \frac{1}{\pi} \int_0^{\infty} dx \left[x \rho(\eta, x) - \frac{2}{1 + \eta x} \right] + \mathcal{O}(s+1). \quad (23)$$

Inserting (23) into the definition (17) we get

$$E_C = \lim_{s \rightarrow -1} \left[\frac{\lambda}{q} \left\{ \frac{1}{\pi \eta} \frac{1}{s+1} + \frac{1}{\pi \eta} \log \tilde{\mu} + \frac{1}{2\pi} \int_0^{\infty} dx \left[x \rho(\eta, x) - \frac{2}{1 + \eta x} \right] - \frac{1}{24} \right\} + \mathcal{O}(s+1) \right], \quad (24)$$

where $\tilde{\mu} = \mu q / \lambda$ is also an arbitrary (due to arbitrariness of μ), but now dimensionless constant.

Because of the pole in the first term the Casimir energy (24) is a divergent quantity and we need to apply a renormalization procedure to remove this divergence.

Let us first notice that the effective action of a single component, free field, which describe the only functional d.o.f. of the string oscillations left after the gauge fixing and imposition of the boundary conditions, has to be of the form:

$$S_{\text{eff}} = - \int_{\tau_1}^{\tau_2} d\tau \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} d\sigma \sqrt{-g^{(0)}} \left\{ \frac{1}{2} g^{(0)ab} \partial_a \psi \partial_b \psi - \frac{1}{2} (m^2 + \xi R^{(0)}) \psi^2 \right\}, \quad (25)$$

where $a, b = \tau, \sigma$,

$$g_{ab}^{(0)} = \frac{q^2}{\lambda^2} \cos^2(\lambda \sigma)$$

is the induced metric tensor on the rotating string worldsheet, $R^{(0)}$ is the induced curvature scalar and ξ, m^2 are constants. Choosing $m = 0, \xi = 1$, we get from (25) the equation of motion which coincides with the equation (8). Moreover, the quantum Hamiltonian which follows from the action functional (25),

$$\hat{H}_{\text{eff}} = \frac{1}{2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} d\sigma \left[\hat{\pi}^2(\tau, \sigma) + \left(\partial_\sigma \hat{\psi}(\tau, \sigma) \right)^2 + V(\sigma) \hat{\psi}^2(\tau, \sigma) \right], \quad (26)$$

differs from the Hamiltonian (15) only by a factor $\frac{q}{\lambda}$ following from the fact that (26) generates translation in the worldsheet time τ and (15) in the laboratory time $X^0 = \frac{q}{\lambda} \tau$, and by a constant term given by the background field energy E_0 (16).

Because the Hamiltonian (26) contains products of quantum fields in the same space-time point, the renormalization procedure has to be applied to it to render its eigenvalues finite. This can be accomplished, [16], by replacing the Hamiltonian $\hat{H}_{\text{eff}}$ by its normal-ordered form $:\hat{H}_{\text{eff}}:$.

The normal ordered field product can be written in the form:

$$:\hat{\psi}^2 := \hat{\psi}^2 + C$$

where C is a counter term. Now, the factor in front of the pole term in (24) is of the form

$$\frac{1}{\pi\eta} = \frac{1}{4\pi} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} d\sigma V(\sigma).$$

This shows that the replacement

$$\hat{\psi}^2 \rightarrow: \hat{\psi}^2 := \hat{\psi}^2 - \frac{1}{2\pi} \frac{1}{s+1} \quad (27)$$

in the Hamiltonian (26) renders its eigenvalues finite. Let us also note that in the limit of flat background the applied procedure is the usual subtraction of the (infinite) constant from the quantum Hamiltonian, which is equivalent to shifting the origin of the energy scale.

After eliminating the divergent term the Casimir energy (24) is still not uniquely determined due to the dependence on the arbitrary parameter $\tilde{\mu}$. This can be of course traced back to the renormalization prescription (27): without spoiling the condition of finiteness of the Hamiltonian eigenvalues we could add to the r.h.s. of (27) arbitrary constant $\log \tilde{\mu}$.

The actual value of $\tilde{\mu}$ cannot be calculated in the presented framework — it is an additional parameter that appears in the model at the quantum level and in order to determine it we need some experimental data. Let us note however that the dependence of the Casimir energy on the quantum scale $\tilde{\mu}$ is, in fact, not surprising. The Casimir energy is intimately related to the one-loop physics [17], and the occurrence of anomalous scale dependence in one-loop field theory is by now a well-understood phenomenon [18, 19]

In order to be able to see the consequences of the formula (24) with divergent term being subtracted let us note that it can be rewritten as a function of the unperturbed string length,

$$L = \frac{2q}{\lambda^2} \sin\left(\frac{\pi\lambda}{2}\right),$$

and a single dimensionless parameter $\kappa = \sqrt{\frac{\alpha}{\gamma L^2}}$:

$$E_C(\kappa) = \frac{2}{\pi L} \left\{ \frac{\sin^2\left(\frac{\pi\lambda(\kappa)}{2}\right)}{\cos\left(\frac{\pi\lambda(\kappa)}{2}\right)} \log \tilde{\mu} + \frac{\sin\left(\frac{\pi\lambda(\kappa)}{2}\right)}{2\lambda(\kappa)} \int_0^\infty dx \left[x\rho(\eta(\kappa), x) - \frac{2}{1 + \eta(\kappa)x} \right] - \frac{1}{24} \right\} \quad (28)$$

where

$$\sin\left(\frac{\pi\lambda(\kappa)}{2}\right) = \sqrt{1 + \kappa^2} - \kappa, \quad \eta(\kappa) = \left[\lambda(\kappa) \tan\left(\frac{\pi\lambda(\kappa)}{2}\right) \right]^{-1},$$

and we have used the relation (5).

The factor in front of the undetermined term $\log \tilde{\mu}$ diverges for $\alpha \rightarrow 0$ (although rather slowly, as $\alpha^{-\frac{1}{4}}$). Consequently, if we demand that our model should give finite value of the Casimir energy in the Nambu–Goto limit we have the unique choice $\log \tilde{\mu} = 0$. The values of the Casimir energy computed in this way, plotted on the Fig.2 below, are negative for all values of the parameters α, γ and for any length L of the unperturbed, rotating string. Consequently, our model does not possess the ground state – we can always lower the energy by taking strings with smaller length L . This result is true for all $\tilde{\mu} < 1$ where the additional term $\sim \log \tilde{\mu}$ lowers the energy further.




Fig 2. The Casimir energy of the NGGB model as a function of the dimensionless parameter

$$\kappa = \sqrt{\frac{\alpha}{\gamma L^2}} \text{ for } \tilde{\mu} = 1.$$

In the Nambu–Goto ($\kappa = 0$) limit formula (28) gives

$$E_C(0) = -\frac{1}{12} \frac{1}{L}.$$

This is different from the celebrated Lüscher term [20],

$$E_C^L = -\frac{\pi}{12} \frac{1}{L},$$

but the reasons are obvious. First, Lüscher term is derived for the string with fixed ends and the oscillation frequencies equal

$$\nu_n^L = \frac{\pi n}{L},$$

while in our, rotating string case we have (14)

$$\nu_n(\kappa = 0) = \frac{2n}{L}.$$

Second, in considered model we have only planar oscillations and this gives additional factor 1/2. This agreement can be viewed as a consistency check for the used regularization method.

If the quantum mechanically generated scale $\tilde{\mu} > 1$, the conclusions concerning the existence of the ground state are different. The first, positive term in the Eq. (28) grows indefinitely for $\alpha \rightarrow 0$ and it is always possible to choose the parameters of the model in such a way that the Casimir energy is positive. Consequently, the semiclassical (i.e. classical + Casimir) string energy,

$$E_{sc} = c_2 L + \frac{c_2}{L},$$

where L is the string length and the positive constants c_1, c_2 can be read off from Eqs. (16) and (28) respectively, possesses positive minimum for some non-zero value of the rotating string length. This configuration can be taken as a correct starting point for the semiclassical quantization procedure.

Acknowledgments.

I would like to thank Prof. Henryk Arodź, Dr Paweł Węgrzyn and Dr Jacek Dziarmaga for numerous helpful discussions. This work is partially supported by the KBN grant 2 P03B 095 13.

References

- [1] J.Polchinski, *Strings and QCD?*, preprint UTTG-92-16 and hep-th/9210045.
- [2] G.'t Hooft, Nucl.Phys **B72** (1974) 461.
- [3] K.Wilson, Phys.Rev **D8** (1974) 2445.
- [4] H.B.Nielsen and P.Olesen, Nucl.Phys. **B61** (1973) 45.
- [5] M.Baker, J.S.Ball and F.Zachariasen, Phys.Rev. **D41** (1990) 2612.

- [6] David J.Gross, Nucl.Phys. **B400** (1993) 161. David J.Gross and Washington Taylor IV, Nucl.Phys. **B400** (1993) 181. David J.Gross and Washington Taylor IV, Nucl.Phys. **B403** (1993) 395.
- [7] Y.Nambu, Lectures on the Copenhagen Summer Symposium (1970), unpublished, O.Hara, Prog.Theor.Phys. **46** (1971) 1549, T.Goto, Prog.Theor.Phys. **46** (1971) 1560, L.N.Chang and J.Mansouri, Phys.Rev. **D5** (1972) 2535, J.Mansouri and Y.Nambu, Phys.Lett. **39B** (1072) 375.
- [8] B.M.Barbashov, V.V.Nesterenko *Introduction to the relativistic string theory*, World Sci., Singapore 1990.
- [9] M.Green, J.Schwarz and E.Witten, *Superstring theory*. Cambridge University Press, Cambridge 1987.
- [10] J.Scherk, Rev.Mod.Phys. **47** (1975) 123.
- [11] H.B.G.Casimir, Proc.Kon.Nederl.Akad.Wetenschap **51** (1948) 739.
- [12] V.V.Nesterenko, Z.Phys. **C51** (1991) 643, H.Kleinert, G.Lambiase and V.V.Nesterenko, Phys.Lett. **B384** (1996) 231, V.V.Nesterenko and I.G.Pirozhenko, Phys.Rev. **D55** (1997) 6603.
- [13] P.Węgrzyn, Phys.Rev. **D50** (1994) 2769.
- [14] L.Hadasz, P.Węgrzyn, Phys.Rev. **D51** (1995) 2891.
- [15] J.Liouville, J.Math.Phys.P.Appl. **18** (1853) 71, A.R.Forshyt, *Theory of Differential Equations*, Part4, vol. 6, New York, Dover, 1959, G.P.Jorjadze, A.K.Pogrebkov, M.C.Polivanov, Doklady Akad. Nauk SSSR **243** (1978) 318.
- [16] R.Rajaraman, *Solitons and Instantons*, Amsterdam, North Holland, 1987.
- [17] S.K.Blau, M.Viser and A.Wipf, Nucl.Phys. **B310** (1988) 163.
- [18] P.Ramond, *Field theory: a modern primer*, Benjamin/Cummings, Reading, 1991.
- [19] C.Itzykson and J. Zuber, *Quantum field theory*, McGraw–Hill, New York, 1980.
- [20] M.Lüscher, Nucl.Phys. **B180** (1981) 317, O.Alvarez, Phys.Rev. **D24** (1981) 440.